Recommended vaccines for adults: 19-65+ years

The chart below shows when you should receive your vaccines. Check the boxes \blacksquare to help you keep track.

Name:					
Vaccine	19-21 years	22-26 years	27-49 years	50-64 years	≥65 years
Chickenpox (varicella)	□ □ 2 doses (if born in 1980 or later)				
Flu (influenza) yearly					
Hepatitis A	□ □ 2 or 3 doses				
Hepatitis B	□ □ 2 or 3 doses				
HIB	□ □ 1 or 3 doses				
HPV	□ □ 1-3 doses (ask your provider)				
Meningococcal MenACWY	☐ ☐ 1 or 2 doses, then booster every 5 years if risk remains				
Meningococcal MenB	□ □ 2 or 3 doses				
MMR (measles, mumps, rubella)	□ □ 1 or 2 doses (if born in 1957 or later)				
Pneumococcal PCV13	□ 1 dose (if born in 1980 or later)				☐ 1 dose
Pneumococcal PPSV23	□ □ 1 or 2 doses				☐ 1 dose
TD or TdaP(tetanus, diphtheria, pertussis)	☐ 1 dose, then Td booster every 10 years (or more as recommended by your provider)				
Zoster-RZ					2 doses
Recommended at age requirement	Recommended for high-risk adults				

You can get this in other languages, large print, braille or a format you prefer. You can also ask for an interpreter. This help is free. Call 855-722-8206 or TTY 711. We accept relay calls.

